

## CORVITE 150

### Supplement Facts

Serving Size: 1 Tablet

Servings Per Container: 30

	Amount per Serving	% Daily Value Adults
<b>Vitamin C</b> (as Calcium Ascorbate)*	120 mg	133%
<b>Vitamin B6</b> (as Pyridoxine HCl)	10 mg	588%
<b>Folate</b> (as Folic Acid 700 mcg and Quatrefolic® ((6S)-5-methyltetrahydrofolic acid) 300 mcg (equivalent to 600 mcg of (6S)-5-methyltetrahydrofolic acid, glucosamine salt)**	1667 mcg DFE	417%
<b>Vitamin B12</b> (as Cyanocobalamin)	20 mcg	833%
<b>Biotin</b>	100 mcg	333%
<b>Iron</b> (as Ferronyl® Carbonyl Iron)	150 mg	833%
<b>Magnesium</b> (as Magnesium Oxide)	10 mg	2%
<b>Zinc</b> (as Zinc Citrate)	25 mg	227%
<b>Calcium Threonate*</b>	5 mg	†

\* (Acid metabolites – such as L-threonic acid, help with absorption of Vitamin C)  
† Daily Value (DV) not established

**Other Ingredients:** Microcrystalline Cellulose, Pregelatinized Starch, di-Calcium Phosphate, Red Color Coating (Hydroxypropylmethyl Cellulose, Polyvinyl Alcohol, FD&C Red #40 Lake, Polyethylene Glycol, Talc, Titanium Dioxide), Croscarmellose Sodium, Stearic Acid, Magnesium Stearate

#### VEGAN FRIENDLY

#### GLUTEN-, LACTOSE- AND SUGAR-FREE

CORVITE®150 is a prescription hematinic multivitamin/multimineral dietary supplement used to improve the nutritional status of patients with iron deficiency

#### CONTRAINDICATIONS

CORVITE®150 should not be used by patients with a known hypersensitivity to any of the listed ingredients. All iron compounds are contraindicated in patients with hemochromatosis, hemosiderosis, or hemolytic anemias.

**WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. KEEP THIS PRODUCT OUT OF REACH OF CHILDREN. In case of accidental overdose, call a doctor or poison control center immediately.**

#### PRECAUTIONS

Do not exceed recommended dose. The type of anemia and the underlying cause or causes should be determined before starting therapy with CORVITE®150. Since the anemia may be a result of a systemic disturbance, such as recurrent blood loss, the underlying cause or causes should be corrected, if possible.

**Folic Acid:** Folic acid alone is an improper therapy in the treatment of pernicious anemia and other megaloblastic anemias where vitamin B<sub>12</sub> is deficient. Folic acid in doses above 0.1 mg daily may obscure pernicious anemia in that hematologic remission can occur while neurologic manifestations remain progressive. Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

The patient's medical conditions and consumption of other drugs, herbs, and/or supplements should be considered.

**Pediatric Use:** Safety and effectiveness in pediatric populations have not been established.

**Geriatric Use:** Safety and effectiveness in elderly populations have not been established.

Biotin levels higher than the recommended daily allowance may cause interference with some laboratory tests, including cardiovascular diagnostic tests (e.g. troponin) and hormone tests, and may lead to incorrect test results. Tell your healthcare provider about all prescription and over-the-counter medicines, vitamins, and dietary supplements that you take, including biotin.

#### DRUG INTERACTIONS

CORVITE® 150 is not recommended for and should not be given to patients receiving levodopa because the action of levodopa is antagonized by pyridoxine. There is a possibility of increased bleeding due to pyridoxine interaction with anticoagulants (e.g., Aspirin, Heparin, Clopidogrel).

#### ADVERSE REACTIONS

Adverse reactions with iron therapy may include constipation, diarrhea, nausea, vomiting, dark stools and abdominal pain. Adverse reactions with iron therapy are usually transient.

#### OVERDOSAGE

The clinical course of acute iron overdosage can be variable. Initial symptoms may include abdominal pain, nausea, vomiting, diarrhea, tarry stools, melena, hematemesis, hypotension, tachycardia, metabolic acidosis, hyperglycemia, dehydration, drowsiness, pallor, cyanosis, lassitude, seizures, shock and coma.

#### DESCRIPTION

CORVITE®150 is a red, oval shaped tablet imprinted with "VP052" on one side and bisected on the other side.

#### DIRECTIONS FOR USE

Take one tablet by mouth daily or as directed by a physician.

#### HOW SUPPLIED

CORVITE®150 is available in bottles of 30 tablets.

**Product Code: 68025-052-30**

#### STORAGE

Store at controlled room temperature 15°-30°C (59°-86°F) [See USP]. Protect from light, moisture and excessive heat. Dispense in a tight, light resistant container as defined in the USP using a child-resistant closure.

#### KEEP THIS PRODUCT OUT OF REACH OF CHILDREN.

\*\*Quatrefolic® is a registered trademark of Gnosis, SpA. U.S. Patent No. 7,947,662.

#### For use on the order of a healthcare practitioner

**Call your doctor about side effects. To report side effects, call Vertical Pharmaceuticals at 1-770-509-4500 or FDA at 1-800-FDA-1088.**

PLR-CORV150-00001-1

Rev. 12/2021

**Distributed by:**  
Vertical Pharmaceuticals, LLC  
Alpharetta, GA 30005  
[www.verticalpharma.com](http://www.verticalpharma.com)

**CORVITE 150**

